

ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN

Allergies/Reactions:

- | | |
|---|--|
| <input type="checkbox"/> Puyallup Infusion Center - Fax: 253-697-5066 | <input type="checkbox"/> Gig Harbor Infusion Services - Fax: 253-530-8069 |
| <input type="checkbox"/> Allenmore Ambulatory Infusion Services - Fax: 253-864-4052 | <input type="checkbox"/> DHEC Infusion Center - Fax: 509-755-5845 |
| <input type="checkbox"/> Auburn Infusion Services - Fax: 253-876-8282 | <input type="checkbox"/> North Spokane Infusion Center - Fax: 509-232-2531 |

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

Ravulizumab-cwvz (Ultomiris)

Patient Name: _____ Requested Date of Service: ____/____/____

Date of Birth: ____/____/____ Patient Phone Number: (____) _____ - _____ May leave message

Diagnosis:

- Atypical Hemolytic Uremic Syndrome (AHUS)
- Myasthenia gravis (MG)
- Paroxysmal Nocturnal Hemoglobinuria (PNH)

ICD -10 Code:

- _____
- _____
- _____

Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs and documentation
****If required documentation not received with order, scheduling of treatment will be delayed until complete information is available****

Baseline labs required: None required

Maintenance labs: None required Other _____ Frequency _____

Baseline Vaccinatin (required): Meningococcal vaccine at least 2 weeks prior to administering initial dose of Ravulizumab-cwvz (Ultomiris)

Date given: ____/____/____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **Patient must be enrolled in Eculizumb (Soliris) REMS program before starting.**

IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P; Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.

Pre-meds: Non recommended

Treatment Regimen:

- 40 to <60 kg = Loading Dose = 2400 mg x 1; Maintenance Dose = 3000 mg q8 weeks starting 2 weeks after LD
- 60 to <100 kg = Loading Dose = 2700 mg x 1; Maintenance Dose = 3300 mg q8 weeks starting 2 weeks after LD
- >= 100 kg = Loading Dose = 3000 mg x 1; Maintenance Dose = 3600 mg q8 weeks starting 2 weeks after LD

Vital Signs: Check vital signs prior to and at completion of infusion.
 Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)

If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.):

- Consult MultiCare Hypersensitivity guideline for treatment/management
- Notify provider of reaction, assessment and need for further orders"

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature

Print Name

Date

Time

Orders expire in 12 months**

Patient Identification - Always Attach Patient Label

Name:
 MRN #:
 CSN #:
 Age / Sex and Gender:

Pre-printed Order
AHUS / MG / PNH



60-0352-3 (5/23)