

ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN

Allergies/Reactions:

- | | |
|---|--|
| <input type="checkbox"/> Puyallup Infusion Center - Fax: 253-697-5066 | <input type="checkbox"/> Gig Harbor Infusion Center - Fax: 253-530-8069 |
| <input type="checkbox"/> Allenmore Ambulatory Infusion Services - Fax: 253-864-4052 | <input type="checkbox"/> DHEC Infusion Center - Fax: 509-755-5845 |
| <input type="checkbox"/> Auburn Infusion Center - Fax: 253-876-8282 | <input type="checkbox"/> MultiCare Adult Home/Alternate Infusion Services - 253-459-6651 |
| <input type="checkbox"/> Tacoma General Infusion Center - Fax: 253-403-1676 | <input type="checkbox"/> North Spokane Infusion Center - Fax: 509-232-2531 |

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

Agalsidase Beta (Fabrazyme)

Patient Name: _____ Requested Date of Service: ____/____/____

Date of Birth: ____/____/____ Patient Phone Number: (____) ____ - ____ May leave message

ICD -10 Code:

- Diagnosis:** Fabry Disease _____
 Other _____ _____

Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs
If required documentation not received with order, scheduling of treatment will be delayed until complete information is available

- IV Access:** Access and/or maintain IV site in accordance with MHS IV Therapy P&P: Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.

Patient weight _____ lb/kg (required)

Treatment Regimen:

- Pre-meds given prior to infusion:
 • Acetaminophen 650 mg po x 1 dose

Agalsidase Beta (Fabrazyme): Administered IV in NS with 0.22 micron filter

Patient wgt <= 35 kg = 50 ml TV; 35.1-70 kg = 100 ml; 70.1-100 kg = 250 ml TV; 100 mg = 500 m TV
 1 mg/kg = _____mg IV every 2 weeks x _____ months (up to 12 months)

Infusion Rate: For patients >= 30 kg: Infusion should be initiated at a rate of 15 mg/hour; after tolerance to initial infusion rate is established, the infusion rate may be increased in increments of 3 to 5 mg/hour with each subsequent infusions. Administration duration: ≥ 1.5 hours (based upon individual tolerability).

- Vital Signs:** Check vital signs prior to and at completion of infusion.
 Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)

If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.):

- Consult MultiCare hypersensitivity guideline for treatment management
- Notify provider of reaction, assessment and need for further orders

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature _____ Print Name _____ Date _____ Time _____

Another brand of drug, identical in form and content, may be dispensed unless checked **Orders expire in 12 months****

Patient Identification - Always Attach Patient Label

Name: _____
 MRN #: _____
 CSN #: _____
 Age / Sex and Gender: _____

Pre-printed Order
AGALSIDASE BETA (Fabrazyme)

