ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN			
Allergies/Reactions:	☐ Puyallup Infusion Center - Fax: 253-697-5066	☐ Gig Harbor Infusion Services - Fo	ax: 253-530-8069
3 .	☐ Allenmore Ambulatory Infusion Services - Fax: 253-86	4-4052 🗖 DHEC Infusion Center - Fax: 509)-755-5845
	☐ Auburn Infusion Services - Fax: 253-876-8282	☐ North Spokane Infusion Center -	Fax: 509-232-2531
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.			
Infliximab (Remicade) or other biosimilars			
Patient Name: Requested Date of Service://			
	Patient Phone Number: ()		
Diagnosis:	ICD -10 Code:	•	5
☐ Rheumatoid Arthritis	<u></u>		
☐ Crohn's Disease ☐ Other	<u> </u>		
Required: H&P with documentation to sup **If required documentation not received v Baseline labs required:	oport above diagnosis including ICD-10 code and with order, scheduling of treatment will be delayed. / Results:		**aldr
 Latent TB testing Date: / 	/ Results:		
HBV screening Date:/	/ Results:		
HCV screening Date:/HIV screening Date:/_	/ Results:		
Maintenance labs required: • Annual Latent TB testing • CBC and CMP annually			
☑ IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P: Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.			
Patient weight = lb/kg (required)			
 Treatment Regimen: □ Pre-meds given 30 minutes prior to infusion (pre-meds recommended for first infusion): Acetaminophen 650 mg po x 1 dose Diphenhydramine 25 mg po x 1 dose or loratadine 10 mg po x 1 dose if intolerant to diphenhydramine 			
☐ Infliximab (Remicade) or other biosimilar (pharmacist to add MHS or insurance preferred product)			
	n between 0.4-4 mg/ml. Attach a 0.22 micron filte		
Once the patient has been established on treatment, MHS will adopt a shortened duration of infusion over 1 hour for patients			
=6 mg/kg. ☐ Yes ☐ No<br Initiation dose:			
0 0	nd to nearest 100 mg) IV at weeks 0, 2, 6 then even		
	nd to nearest 100 mg) IV at weeks 0, 2, 6 then even		
mg/kg = mg (round to nearest 100 mg) IV everyweeks Maintenance dose:			
□ Continue maintenance dose of mg/kg (round to nearest 100 mg) IV everyweeks □ 3 mg/kg =mg (round to nearest 100 mg) IV at weeks 0, 2, 6 then every 8 weeks □ 5 mg/kg =mg (round to nearest 100 mg) IV at weeks 0, 2, 6 then every 8 weeks □mg/kg =mg (round to nearest 100 mg) IV everyweeks □ Continue maintenance dose ofmg (round to nearest 100 mg) IV everyweeks			
☑ Vital Signs: Check vital signs prior to and at completion of infusion. Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)			
 If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.): Consult MultiCare Hypersensitivity guideline for treatment/management Notify provider of reaction, assessment and need for further orders" 			
Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.			
Was consent obtained: ☐ Yes ☐ No (if yes, please send DOCUMENTATION of consent with order)			
Provider Signature	Print Name	Date Time	
-	content, may be dispensed unless checked 🗖	Orders expire in 12 mo	nths**
Patient Identification - Always Attach Patient Label Pre-printed Order			

Name:

MRN #:

CSN #:

Age / Sex and Gender:

Pre-printed Order
INFLIXIMAB (Remicade)
and Biosimilars

MultiCare 🕰

