ALL ORDERS MUST BE SIGNED AND DATED BY THE REFERRING PROVIDER	
ALLERGIES/REACTIONS (REQUIRED):	Yakima Outpatient Infusion Care 808 N 39 th Ave Yakima WA 98902
	Phone: 509-575-1174
	Fax: 509-577-5021
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.	
CODE STATUS Patients will be considered FULL CODE unless marked otherwise. If the patient has a POLST, advance directive, or living will, please include a copy with the orders.	
ABATACEPT (Orencia)	
Datient Name:	Requested Start Date:/
Patient Name:	kg Patient Height:
DIAGNOSIS & ICD-10 CODE:	
☐ Rheumatoid Arthritis (ICD-10:) ☐ Other:	(ICD-10:)
REQUIRED: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs **If required documentation not received with this order, scheduling of treatment will be delayed until complete information is available **	
REQUIRED BASELINE LABS & INFORMATION: ✓ CBC & CMP	
✓ CBC & CMP ✓ Negative Latent TB Test (Date: □ QuantiFERON Gold □ PPD □ Chest X-Ray □ Other:)	
✓ HBV Screening/Status (Date: HepBsAg Hep	DBSAb HepB Core Ab)
G. ,, , , <u></u> , , , , <u></u> ,	
ROUTINE LABS: ☐ CMP ☐ CBC w/ diff ☐ LFT ☐ CRP ☐ ESR ☐ Other:	
ROUTINE LAB FREQUENCY: ☐ Each Infusion ☐ Annually ☐ Other:	
ACCESS: Access and maintain IV site or Port-A-Cath in accordance with the appropriate MYM OIC P&Ps	
TREATMENT REGIMEN: (another brand of drug, identical in form and content may be dispensed unless "DAW" or "BRAND ONLY" is written next to the drug name)	
PRE-MEDICATIONS (PRN) – Give 30 minutes prior to infusion ☐ Acetaminophen 650 mg PO x1 ☐ Diphenhydramine 25 mg PO x1 ☐ Diphenhydramine 50 mg IV x1	
☐ Loratadine 10 mg PO x1 ☐ Methylprednisolone sodium succinate 125 mg IV x1	
□ Other:	
ABATACEPT (Orencia) in 100 mL 0.9% sodium chloride IV infusion over 30 minutes	
☐ Weight < 60 kg = 500 mg	
 ☐ Weight 60 kg to 100 kg = 750 mg ☐ Weight > 100 kg = 1,000 mg 	
☐ Other dose & frequency:	
FREQUENCY: INDUCTION: Infuse every 2 weeks x 3 doses then every 4 weeks thereafter	
☐ MAINTENANCE: Infuse every 4 weeks	
MAINTENANCE. IIIuse every 4 weeks	
MONITORING: Vitals at baseline and at completion of infusion. SUPPORTIVE CARE: Administer hypersensitivity reaction/anaphylaxis management per MYMH OIC protocol as necessary.	
DISCHARGE: 30 minutes after infusion is complete when vital signs are stable and hypersensitivity symptoms are absent. Waiting period can be waived by patient on subsequent infusions.	
Provider Signature:	Date:
Print name:P	hone # Fax #
□ NEW REFERRAL □ UPDATED REFERRAL	**Expires 12 months from written date**
	Expires 12 months from written date
Patient Identification - Attach Patient Label ABATACEPT (Orencia)	

Name:

MRN:

Age / Sex and Gender:

MultiCare 🞜 Yakima Memorial Hospital