

ALL ORDERS MUST BE SIGNED AND DATED BY THE REFERRING PROVIDER

ALLERGIES/REACTIONS (REQUIRED):

Yakima Outpatient Infusion Care
808 N 39th Ave Yakima WA 98902
Phone: 509-575-1174
Fax: 509-577-5021

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

CODE STATUS

Patients will be considered FULL CODE unless marked otherwise. If the patient has a POLST, advance directive, or living will, please include a copy with the orders.

ALPHA₁-PROTEINASE INHIBITOR [HUMAN] (i.e., Prolastin)

Patient Name: _____ Requested Start Date: ____/____/____
Date of Birth: ____/____/____ Patient Weight: _____ kg Patient Height: _____

DIAGNOSIS & ICD-10 CODE:

- Alpha₁-Antitrypsin Deficiency (ICD-10: _____)
- Other: _____ (ICD-10: _____)

REQUIRED: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs

****If required documentation not received with this order, scheduling of treatment will be delayed until complete information is available****

ADDITIONAL INFORMATION / DIRECT DISTRIBUTION NOTE:

- ✓ Additional information may be required by the drug manufacturer's direct distribution program prior to initial and continuation referrals.

ACCESS: Access and maintain IV site or Port-A-Cath in accordance with the appropriate MYM OIC P&Ps

TREATMENT REGIMEN: (another brand of drug, identical in form and content may be dispensed unless "DAW" or "BRAND ONLY" is written next to the drug name)

ALPHA₁-PROTEINASE INHIBITOR [HUMAN] PRODUCT FOR INTRAVENOUS INFUSION

- Prolastin-C Liquid (preferred/formulary)
- Other: _____

DOSAGE & FREQUENCY

- Standard: 60 mg/kg x _____ kg = _____ mg (+/- 10%) IV weekly x 1 year
- Other: _____ mg/kg x _____ kg = _____ mg (+/- 10%) IV _____ (frequency) x _____ (duration)

INFUSE AS TOLERATED BY PATIENT IN ACCORDANCE WITH PRODUCT-SPECIFIC PRESCRIBING INFORMATION

MONITORING: Vitals at baseline, every 15 minutes during, and at completion of infusion.

SUPPORTIVE CARE: Administer hypersensitivity reaction/anaphylaxis management per MYMH OIC protocol as necessary.

DISCHARGE: 30 minutes after infusion is complete when vital signs are stable and hypersensitivity symptoms are absent. Waiting period can be waived by patient on subsequent infusions.

Provider Signature: _____ Date: _____
Print name: _____ Phone # _____ Fax # _____

- NEW REFERRAL
- UPDATED REFERRAL

****Expires 12 months from written date****

Patient Identification - Attach Patient Label

Name: _____

MRN: _____

Age / Sex and Gender: _____

ALPHA₁-PROTEINASE INHIBITORS (i.e., Prolastin)

MultiCare 

Yakima Memorial Hospital