MHS Home/Alternate Site Infusion Services	
Abatacept (Orencia) Infusion Order Set	
ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER	
Patient Name:	DOB:/ Weight: lb/kg
Patient Phone Number: ()	Requested Date of Service:/
Patient Allergies:	
Diagnosis: ICD – 10 Code: • Z45.2: Encounter for adjustment and management of vascular access device • Z95.828: Presence of other vascular implants and grafts	
Baseline labs required: • Latent TB Testing/Quantiferon: Date:/ Result: • HBV Screening Date:/ Result: Maintenance labs required: • Latent TB Testing/Quantiferon every 12 months	
Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. ** Emergency phone number for provider (required) **	
Provider Order for Abatacept (Orencia) Infusion	
l · · · · · · · · · · · · · · · · · · ·	e with check boxes), providers are responsible for indicating a se order. Orders left unchecked will not be initiated.
Medication: Abatacept (Orencia) in 100 mL NS IV infusion over 30 minutes with 0.2 micron filter Patient weight less than 60 kg; Dose = 500 mg Patient weight 60-100 kg; Dose = 750 mg Patient weight greater than 100 kg; Dose = 1000 mg Initiation dose: Every 2 weeks x 3 doses, then every 4 weeks for maintenance dose Maintenance dose: Every 4 weeks Additional Medications for vascular access maintenance: 0.99% NaCl (NS) PF flush in 10 mL syringe: Administer 5 – 30 mL IV as needed for establishing and maintaining IV access and for flushing IV line post infusion. Heparin: 100 units/mL for port OR 10 units/mL for other central line/midline, 3-5 mL PRN for de-accessing line. Cathflo (alteplase) 2 mg: Instill 2 mg (2 mL) into occluded catheter as needed for CVC/CVAD occlusion. Skilled Nurse to draw maintenance labs as follows: Latent TB Testing/Quantiferon every 12 months Other Lab(s): Skilled Nurse Interventions: Admit (first visit) patient to services for home/alternate site infusion therapy of Abatacept (Orencia). Complete Skilled Nurse visit with each infusion for ongoing home/alternate site infusion therapy of Abatacept (Orencia). Obtain vital signs (TPR & B/P) at baseline and at completion of infusion. Obtain patient weight at each visit. Establish IV access and flush per policy to maintain patency. CONTINUED ON NEXT PAGE	
FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY	
Patient Identification: Name: MRN: DOB:	Pre-printed order – Page 1 of 2 Abatacept (Orencia) MultiCare

- Reconstitute each vial of Abatacept with 10 mL sterile water using the supplied silicone-free syringes only and an 18- to 21-gauge needle. DO NOT SHAKE.
- Add reconstituted medication to sodium chloride 0.9% (NS) for a total volume of 100 mL.
- Infuse Abatacept (Orencia) as prescribed.
- Once infusion complete, flush IV line with sodium chloride 0.9% (NS) as prescribed.
- If vital signs are stable after infusion, Skilled Nurse will discontinue IV access and complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

MRN:

DOB:

- STOP THE INFUSION.
- Administer emergency medications as prescribed (below).
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- Diphenhydramine 50 mg/mL solution: Inject 0.5 mL (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available.
- Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution: Inject 2 mL (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- Acetaminophen (Tylenol) 325 mg tab: Take 2 tablets (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- 0.9% NaCl (NS) solution: Infuse 500 mL into the vein via gravity as needed at a rate needed to maintain IV access.
- Oxygen: Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders. Was consent obtained:

Yes

No (if yes, please send DOCUMENTATION of consent with order) I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services. Provider Signature: _____ ____ Printed Name: __ ___ Orders expire in 12 months unless otherwise specified: ___ Provider/Clinic Information: Return completed orders to: MultiCare Home/Alternate Infusion Services Address: ___ 253-459-6650 (phone) / 253-864-2785 (fax) Phone #: _____ Fax#: ____ FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY Patient Identification: Pre-printed order – Page 2 of 2 Name: Abatacept (Orencia)

MultiCare

Revised 06/24