

MHS Home/Alternate Site Infusion Services Iron Infusion Order Set

ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER

Patient Name: _____	DOB: ____/____/____	Weight: _____ lb/kg
----------------------------	----------------------------	----------------------------

Patient Phone Number: (____) _____-_____	Requested Date of Service: ____/____/____
---	--

Patient Allergies: _____

Diagnosis: _____ **ICD – 10 Code:** _____

- Z45.2: Encounter for adjustment and management of vascular access device
- Z95.828: Presence of other vascular implants and grafts

Baseline Labs (Required):

- | | | |
|--------------------------|----------------------|---------------|
| • Hgb | Date: ____/____/____ | Result: _____ |
| • Serum Creatinine | Date: ____/____/____ | Result: _____ |
| • Ferritin | Date: ____/____/____ | Result: _____ |
| • Iron | Date: ____/____/____ | Result: _____ |
| • Transferrin saturation | Date: ____/____/____ | Result: _____ |

Parenteral iron products are indicated for patients who have intolerance to or failed oral iron. **Was oral iron tried?**

- Yes No Medical Necessity: _____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **** Emergency phone number for provider _____ (required) ****

Provider Order for Iron Infusion

ORDERS WITH CHECK
BOXES

When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

Iron Infusion:

- Ferumoxytol (Feraheme) 510 mg IV infusion over 20 minutes every week for 2 doses (diluted in 100 mL NS)
- Ferric Carboxymaltose (Injectafer) 750 mg IVP (100 mg per minute) every week for 2 doses (undiluted)
- Iron Sucrose (Venofer):
 - 100 mg infused over at least 10 minutes every 48 hours for _____ doses (undiluted)
 - 200 mg infused over at least 15 minutes every 48 hours for _____ doses (undiluted)
 - 300 mg infused over at least 90 minutes weekly for _____ doses (diluted in 150 mL NS)
 - Other: _____
- Other: _____

Additional Medications for vascular access maintenance:

- 0.9% NaCl (NS) PF flush in 10 mL syringe: Administer 5 – 30 mL IV as needed for establishing and maintaining IV access and for flushing IV line post infusion.
- Heparin: 100 units/mL for port OR 10 units/mL for other central line/midline, 3-5 mL PRN for de-accessing line.
- Cathflo (alteplase) 2 mg: Instill 2 mg (2 mL) into occluded catheter as needed for CVC/CVAD occlusion.

Skilled Nurse to draw labs as follows:

Lab(s): _____ Frequency: _____

CONTINUED ON NEXT PAGE

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:
MRN:
DOB:

Pre-printed order – Page 1 of 2
Iron Order Set
MultiCare

Skilled Nurse Interventions:

- Admit (first visit) patient to services for home infusion therapy of Iron.
- Complete Skilled Nurse Visit with each infusion for ongoing home infusion therapy of Iron.
- Obtain vital signs (TPR & B/P) at baseline, at completion of infusion, and 30 minutes post-infusion.
- Obtain patient weight at each visit.
- Establish IV access and flush per policy to maintain patency.
- Draw labs as ordered.
- Add Iron to required volume of sodium chloride 0.9% (NS) or give undiluted, depending on product selected.
- Infuse Iron as prescribed.
- Once infusion complete, flush IV line with sodium chloride 0.9% (NS) as prescribed.
- If vital signs are stable 30 minutes after infusion, Skilled Nurse will discontinue IV access and complete visit. If no infusion-related events with previous 3 infusions, may waive post-infusion monitoring and discharge patient home at completion of infusion.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

- STOP THE INFUSION.
- Administer emergency medications as prescribed (below).
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available.
- **Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution:** Inject **2 mL** (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- **Acetaminophen (Tylenol) 325 mg tab:** Take **2 tablets** (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- **0.9% NaCl (NS) solution:** Infuse **500 mL** into the vein via gravity as needed at a rate needed to maintain IV access.
- **Oxygen:** Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.

Provider Signature: _____ Printed Name: _____ Date: _____

NPI: _____ Orders expire in 12 months unless otherwise specified: _____

Provider/Clinic Information:

Address: _____

Phone #: _____ Fax#: _____

Return completed orders to:

MultiCare Home/Alternate Infusion Services
253-459-6650 (phone) / 253-864-2785 (fax)

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:
MRN:
DOB:

Pre-printed order – Page 2 of 2
Iron Order Set
MultiCare

Revised 06/24