MHS Home/Alternate Site Infusion Services Pembrolizumab (Keytruda) Infusion Order Set ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER DOB: ____/___ **Patient Name: Weight:** _____ lb/kg Patient Phone Number: (_____) ____-Requested Date of Service: ____/___ **Patient Allergies:** Diagnosis: _____ ICD – 10 Code: Z45.2: Encounter for adjustment and management of vascular access device Z95.828: Presence of other vascular implants and grafts Baseline Labs required: CBC/CMP Result: _____ Date: ____/____ HBV screening Date: ____/___ Result: _____ Date: ____/___ Result: _____ HCV screening Date: / /___ Result: ____ HIV screening TSH/Cortisol Result: Maintenance Labs: Monitor thyroid function, cortisol, CBC, and CMP periodically during treatment. Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. ** Emergency phone number for provider ______ (required) ** Provider Order for Pembrolizumab (Keytruda) Infusion ORDERS WITH CHECK When an order is optional (those with check boxes), providers are responsible for indicating a BOXES check mark in the box next to the order. Orders left unchecked will not be initiated. **Biologic Infusion:** Pembrolizumab (Keytruda) IV added to 100 mL NS. Infuse over 30 minutes via infusion pump using 0.22 micron in-line filter □ 200 mg IV every 3 weeks □ 400 mg IV every 6 weeks Dosing Schedule: Give dose ______ to ____ days BEFORE chemotherapy. ☐ Give dose _____ to ____ days AFTER chemotherapy. **Additional Medications** for vascular access maintenance: 0.9% NaCl (NS) PF flush in 10 mL syringe: Administer 5 – 30 mL IV as needed for establishing and maintaining IV access and for flushing IV line post infusion. Heparin: 100 units/mL for port OR 10 units/mL for other central line/midline, 3-5 mL PRN for de-accessing line. Cathflo (alteplase) 2 mg: Instill 2 mg (2 mL) into occluded catheter as needed for CVC/CVAD occlusion. Skilled Nurse to draw labs as follows: CBC w/ diff and CMP every 3 weeks Or other frequency: _____ □ TSH/Cortisol every 3 weeks Or other frequency: ____ Skilled Nurse Interventions: Admit (first visit) patient to services for home infusion therapy of Pembrolizumab (Keytruda). Complete Skilled Nurse visit with each infusion for ongoing home infusion therapy of Pembrolizumab (Keytruda). Obtain vital signs (TPR & B/P) at baseline and at completion of infusion. CONTINUED ON NEXT PAGE FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY Patient Identification: Pre-printed order – Page 1 of 2 Pembrolizumab (Keytruda) Order Set Name: MRN: MultiCare DOB:

- Obtain patient weight at each visit.
- Establish IV access and flush per policy to maintain patency.
- Draw labs as ordered.
- Add Pembrolizumab (Keytruda) to 100 mL of sodium chloride 0.9% (NS).
- Infuse Pembrolizumab (Keytruda) as prescribed.
- Once infusion complete, flush IV line with sodium chloride 0.9% (NS) as prescribed.
- If vital signs are stable after completion of infusion, Skilled Nurse will discontinue IV access and complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

- STOP THE INFUSION.
- Administer emergency medications as prescribed (below).
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available.
- Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution: Inject 2 mL (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- Acetaminophen (Tylenol) 325 mg tab: Take 2 tablets (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- 0.9% NaCl (NS) solution: Infuse 500 mL into the vein via gravity as needed at a rate needed to maintain IV access.
- Oxygen: Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained:

Yes
No (if yes, please send DOCUMENTATION of consent with order)

I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.

Provider Signature:
Printed Name:
Date:
NPI:
Orders expire in 12 months unless otherwise specified:
MultiCare Home/Alternate Infusion Services 253-459-6650 (phone) / 253-253-864-2785 (fax)

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

_____Fax#: __

Name:

Phone #: ____

MRN: DOB: Pre-printed order – Page 2 of 2

Pembrolizumab (Keytruda) Order Set

MultiCare Revised 05/24