ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN				
Allergies/Reactions:	☐ Puyallup Infusion Center - Fax: 25	3-697-5066	☐ Gig Harbo	r Infusion Services - Fax: 253-530-8069
	☐ Allenmore Infusion Services - Fax	253-864-4052	☐ DHEC Infu	sion Center - Fax: 509-755-5845
	☐ Auburn Infusion Services - Fax: 25		·	kane Infusion Center - Fax: 509-232-2531
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.				
Risankizumab (Skyrizi)				
Patient Name:Requested Date of Service:/				
Date of Birth:/	Patient Phone Number: (🗖 May leave message
		ICD -10 Code:		
Diagnosis: □ Crohn's Disease		_		
☐ Ulcertive Colitis		_		
Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs and documentation. **If required documentation not received with order, scheduling of treatment will be delayed until complete information is available**				
Baseline labs required: CBC, CMP Latent TB testing, Date:/ Results: HBV screening, Date:/ Results: HCV screening, Date:/ Results: HIV screening, Date:/ Results: *Patients should be up to date with all immunizations before initiating therapy. Avoid the use of live vaccines in patients undergoing Risankizumab treatment				
☑ IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P: Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.				
Treatment Regimen for Risankizumab (Skyrizi) dose: ☐ Crohn's Disease - 600 mg IV infusion over 1 hour at week 0, 4, and 8. ☐ Ulcerative Colitis - 1200 g IV over 2 hours at week 0, 4, and 8. Maintenance dose for both: 180 pr 360 mg SUBQ at week 12 then every 8 weeks thereafter. **Medicare will not cover SUBQ in outpatient setting				
☑ Vital Signs: Check vital signs prior to and at compeltion of infusion.				
Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)				
If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.): • Consult MultiCare hypersensitivity guideline for treatment management • Notify provider of reaction, assessment and need for futher orders				
Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.				
Provider Signature	Print Name		Date	Time
				Order expires in 12 months**

Patient Identification - Always Attach Patient Label

Name:

MRN #:

CSN #:

Pre-printed Order CROHN'S DISEASE MultiCare



Age / Sex and Gender: