

**ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN**

**Allergies/Reactions:**

Fax all infusions to: 833-380-8800  
Please mark the appropriate infusion center:

- Allenmore Infusion Center
- Auburn Infusion Center
- Gig Harbor Infusion Services
- Puyallup Infusion Center
- DHEC Infusion Center
- North Spokane Infusion Center
- North Star Lodge Infusion Center

**ORDERS WITH CHECK BOXES**

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

**Burosumab-twza (Crysvita):**

Patient Name: \_\_\_\_\_ Requested Date of Service: \_\_\_\_/\_\_\_\_/\_\_\_\_  
Date of Birth: \_\_\_\_/\_\_\_\_/\_\_\_\_ Patient Phone Number: (\_\_\_\_) \_\_\_\_\_ - \_\_\_\_\_  May leave message

**ICD -10 Code:**

**Diagnosis:**  Hypophosphatemia, X-linked  \_\_\_\_\_

**Required:** H&P with documentation to support above diagnosis including ICD-10 code and supporting labs  
**\*\*If required documentation not received with order, scheduling of treatment will be delayed until complete information is available\*\***

**Baseline Labs Required:**

- Fasting serum phosphorus level prior to first dose: Date of test: \_\_\_\_\_ Phosphorus level \_\_\_\_\_  
\*\* fasting serum phosphorus concentration should be below the reference range (2.5-4.5 mg/dL) prior to initiation of treatment

**Maintenance labs required:**

- Fasting serum phosphorus level 2 weeks post-dose then every month for the first 3 months of treatment.  
Verify level prior to giving

**Treatment Regimens:**

- Burosumab-twza (Crysvita) given SUBQ
- 1 mg/kg = \_\_\_\_\_ mg every 4 weeks. Maximum dose is 90 mg. Round to the nearest 10 mg.

**Dose Adjustments:** Serum phosphorus above normal range, hold next dose; reassess fasting serum every 4 weeks; once serum phosphorus falls below the normal range, may reinstate burosumab-twza at a reduced dose (approx. half the initial starting dose). Recheck fasting serum phosphorus every 2 weeks after dose adjustment; based on results, determine if additional dosing adjustment is necessary.

- Vital signs:** Check vital signs prior to and at completion of dose.  
Contact provider if systolic BP>180; diastolic BP>100; systolic BP<90; HR >110; temp >38C (100.4F)

**If Hypersensitivity reaction (fever, chills, hypotension, rigors, itching, rash, etc.)**

- Consult MultiCare hypersensitivity guideline for treatment management
- Notify provider of reaction, assessment and need for further orders

**Code Status:** Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

**Was consent obtained:**  Yes  No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature \_\_\_\_\_ Print Name \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Another brand of drug, identical in form and content, may be dispensed unless checked  **Orders expires in 12 months\*\***

**Patient Identification - Always Attach Patient Label**

Name:  
MRN #:  
CSN #:  
Age / Sex and Gender:

Pre-printed Order  
**HYPOPHOSPHATEMIA**

