

**ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN**

**Allergies/Reactions:**

Fax all infusions to: 833-380-8800  
Please mark the appropriate infusion center:

- Allenmore Infusion Center
- Auburn Infusion Center
- Gig Harbor Infusion Services
- Puyallup Infusion Center
- DHEC Infusion Center
- North Spokane Infusion Center
- North Star Lodge Infusion Center

**ORDERS WITH CHECK BOXES**

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

**Omalizumab (Xolair)**

Patient Name: \_\_\_\_\_ Requested Date of Service: \_\_\_\_/\_\_\_\_/\_\_\_\_

Date of Birth: \_\_\_\_/\_\_\_\_/\_\_\_\_ Patient Phone Number: (\_\_\_\_) \_\_\_\_-\_\_\_\_  May leave message

**ICD -10 Code:**

- Diagnosis:**
- Allergic Asthma (see lab requirement)  \_\_\_\_\_
  - Chronic Idiopathic Urticaria  \_\_\_\_\_
  - Other \_\_\_\_\_  \_\_\_\_\_

**Required:** H&P with documentation to support above diagnosis including ICD-10 code and supporting labs  
*\*\*If required documentation not received with order, scheduling of treatment will be delayed until complete information is available\*\**

**Baseline lab required for initial dosing for diagnosis of allergic asthma:**

- IgE level Date \_\_\_\_\_ Results \_\_\_\_\_

Patient weight = \_\_\_\_\_lb/kg required for allergic asthma diagnosis (not needed for chronic idiopathic urticaria)

*Patient must carry an epinephrine auto-injector in the event of anaphylaxis*

**Treatment Regimen:**

Omalizumab (Xolair) given SQ:

- 150 mg SQ  every 2 weeks or  every 4 weeks
- 300 mg SQ  every 2 weeks or  every 4 weeks
- Other \_\_\_\_\_ mg SQ  every 2 weeks or  every 4 weeks

**Vital Signs:** Check vital signs prior to and after injection.  
Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)

**Special instructions:** If stable 30 minutes post injection, may discharge home. If no injection-related events with previous 3 doses may waive post-injection monitoring period and discharge patient home after completion.

**If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.):**

- Consult MultiCare hypersensitivity guideline for treatment management
- Notify provider of reaction, assessment and need for further orders

**Code Status:** Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

**Was consent obtained:**  Yes  No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature \_\_\_\_\_ Print Name \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Another brand of drug, identical in form and content, may be dispensed unless checked  **Orders expire in 12 months\*\***

**Patient Identification - Always Attach Patient Label**

Name: \_\_\_\_\_  
MRN #: \_\_\_\_\_  
CSN #: \_\_\_\_\_  
Age / Sex and Gender: \_\_\_\_\_

Pre-printed Order  
**OMALIZUMAB (Xolair)**

