

ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN

Allergies/Reactions:

Fax all infusions to: 833-380-8800
Please mark the appropriate infusion center:

- | | |
|---|---|
| <input type="checkbox"/> Allenmore Infusion Center | <input type="checkbox"/> DHEC Infusion Center |
| <input type="checkbox"/> Auburn Infusion Center | <input type="checkbox"/> North Spokane Infusion Center |
| <input type="checkbox"/> Gig Harbor Infusion Services | <input type="checkbox"/> North Star Lodge Infusion Center |
| <input type="checkbox"/> Puyallup Infusion Center | |

ORDERS WITH CHECK BOXES

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

Eculizumab (Soliris)

Patient Name: _____ Requested Date of Service: ____/____/____

Date of Birth: ____/____/____ Patient Phone Number: (____) _____ - _____ May leave message

Diagnosis:

- Atypical Hemolytic Uremic Syndrome (AHUS)
- Myasthenia gravis (MG)
- Neuromyelitis Optica Spectrum Disorder (NOSD)
- Paroxysmal Nocturnal Hemoglobinuria (PNH)

ICD -10 Code:

- _____
- _____
- _____
- _____

Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs and documentation
If required documentation not received with order, scheduling of treatment will be delayed until complete information is available

Baseline labs required: None required

Maintenance labs: None required

Baseline Vaccinatin (required): Meningococcal vaccine at least 2 weeks prior to administering initial dose of Eculizumab (Soliris)

Date given: ____/____/____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **Patient must be enrolled in Eculizumb (Soliris) REMS program before starting.**

IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P; Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.

Pre-meds: Non recommended

Treatment Regimen:

- AHUS/MG/NOSD = 900 mg weekly x 4 doses; then 1200 mg at week 5 then 1200 mg every 2 weeks
- PNH = 600 mg weekly x 4 doses; then 900 mg at week 5 then 900 mg every 2 weeks

Vital Signs: Check vital signs prior to and at completion of infusion.
Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)

If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.):

- Consult MultiCare Hypersensitivity guideline for treatment/management
- Notify provider of reaction, assessment and need for further orders"

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature

Print Name

Date

Time

Orders expire in 12 months**

Patient Identification - Always Attach Patient Label

Name:
MRN #:
CSN #:
Age / Sex and Gender:

Pre-printed Order

AHUS / MG / NOSD / PNH

MultiCare 



60-0350-1 (Rev. 9/24)