

**ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN**

**Allergies/Reactions:**

Fax all infusions to: 833-380-8800  
Please mark the appropriate infusion center:

- Allenmore Infusion Center
- Auburn Infusion Center
- Gig Harbor Infusion Services
- Puyallup Infusion Center
- DHEC Infusion Center
- North Spokane Infusion Center
- North Star Lodge Infusion Center

**ORDERS WITH CHECK BOXES**

When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.

**Pegloticase (Krystexxa):**

Patient Name: \_\_\_\_\_ Requested Date of Service: \_\_\_\_/\_\_\_\_/\_\_\_\_

Date of Birth: \_\_\_\_/\_\_\_\_/\_\_\_\_ Patient Phone Number: (\_\_\_\_) \_\_\_\_\_ - \_\_\_\_\_  May leave message

**ICD -10 Code:**

**Diagnosis:**  Gout  \_\_\_\_\_

**Required:** H&P with documentation to support above diagnosis including ICD-10 code and supporting labs  
*\*\*If required documentation not received with order, scheduling of treatment will be delayed until complete information is available\*\**

**Baseline Labs Required:**

- Screen patients at risk for G6PD deficiency prior to starting pegloticase. Hemolysis and methemoglobinemia have been reported in patients with G6PD deficiency.
- Serum uric acid levels

**Maintenance labs required:**

- Serum uric acid levels prior to infusions, drawn 24-48 hours before the infusion

**Treatment Regimens:**

Pegloticase (Krystexxa)

- 8 mg IV infusion every 2 weeks as monotherapy or co-administered with weekly oral methotrexate and folic acid supplementation; begin methotrexate and folic acid at least 4 weeks prior to starting pegloticase.

Premeds:  Diphenhydramine 25 mg IVP  Methylprednisolone 40 mg IVP

**Dose Adjustments:** Discontinue treatment if levels increase to above 6 mg/dL, particularly when 2 consecutive levels above 6 mg/dL are observed.

- Vital signs:** Check vital signs prior to and at completion of dose.  
Contact provider if systolic BP>180; diastolic BP>100; systolic BP<90; HR >110; temp >38C (100.4F)

**If Hypersensitivity reaction (fever, chills, hypotension, rigors, itching, rash, etc.)**

- Consult MultiCare hypersensitivity guideline for treatment management
- Notify provider of reaction, assessment and need for further orders

**Code Status:** Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

**Was consent obtained:**  Yes  No (if yes, please send DOCUMENTATION of consent with order)

Provider Signature \_\_\_\_\_ Print Name \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Another brand of drug, identical in form and content, may be dispensed unless checked  **Orders expires in 12 months\*\***

**Patient Identification - Always Attach Patient Label**

Name:  
MRN #:  
CSN #:  
Age / Sex and Gender:

Pre-printed Order  
**GOUT**

**MultiCare** 

