ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN			
Allergies/Reactions:	Fax all infusions to: 833-380-8800 Please mark the appropriate infusion center:	☐ Allenmore Infusion Center☐ Auburn Infusion Center☐ Gig Harbor Infusion Services☐ Puyallup Infusion Center	□ DHEC Infusion Center □ North Spokane Infusion Center □ North Star Lodge Infusion Center
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.			
Tocilizumab (Actemra) or other biosimilar			
Patient Name: Requested Date of Service://			
Date of Birth:/Pati	ent Phone Number: () _		🗖 May leave message
ICD -10 Code:			
Diagnosis: ☐ Rheumatoid Arthritis			
☐ Other			
Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs **If required documentation not received with order, scheduling of treatment will be delayed until complete information is available.**			
Baseline labs required for initial dosing: • CBC & CMP (do not initiate if ANC <2000; platelets <100k, and/or liver enzymes > 1.5 x ULN) • Latent TB testing			
DateResults			
Maintenance Labs required: • CBC & CMP every 8 weeks - Hold infusion and notify provider for ANC <1000; platelets <100k; and/or liver enzymes >1.5 x ULN • Lipid panel at 8 weeks, then every 6 months • Annual latent TB testing			
☑ IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P, Peripheral IV device site selection, insertion, maintenance, and discontinuation and maintenance of central venous catheters-flushing, dressing changes and removal.			
Patient weight =	lb/kg (require	ed)	
TREATMENT REGIMEN (pharmacist to add MHS or insurance preferred product): Tocilizumab or other biosimilar: administered in 100 mL NS infused over 60 minutes			
\square 4 mg/kg =mg (maximum dose = 800 mg) IV every 4 weeks			
■ 8 mg/kg = mg (maximum dose = 800 mg) IV every 4 weeks			
☑ Vital Signs: Check vital signs prior to and after infusion. Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)			
If Hypersensitivity reaction (fever, chills, hypotension, rigors, itching, rash, etc.): • Consult MultiCare hypersensitivity guideline for treatment management • Notify provider of reaction, assessment and need for further orders			
Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.			
Was consent obtained: ☐ Yes ☐ No (if yes, please send DOCUMENTATION of consent with order)			
Provider Signature	Print Name	Date	Time
Another brand of drug, identical in form and content,	may be dispensed unless checked		Order expires in 12 months**
Patient Identification - Always Attach Patient Label			

Name:

MRN #:

CSN #:

Age / Sex and Gender:

Pre-printed Order
TOCILIZUMAB (Actemra)

MultiCare 🕰

