ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN			
Allergies/Reactions:	Fax all infusions to: 833-380-8800 Please mark the appropriate infusion center:	☐ Allenmore Infusion Center☐ Auburn Infusion Center☐ Gig Harbor Infusion Services☐ Puyallup Infusion Center☐	□ DHEC Infusion Center □ North Spokane Infusion Center □ North Star Lodge Infusion Center
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.			
Osteoporosis Treatments			
Patient Name:Requested Date of Service://			
		May leave message	
Diagnosis: ☐ Osteoporosis ☐ Osteopenia ☐ Other		<u>Code:</u>	
Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs. Dexa scan is recommended every			
2 years. **If required documentation not received with order, scheduling of treatment will be delayed until complete information is available**			
Baseline labs required: • BMP • Serum Calcium mg/dL Contraindicated to give denosumab, romosozumab-aqqg or bisphosphanates in patients with hypocalcemia • Serum Creatinine mg/dL Contraindicated to give zoledronic acid if CrCl <35 mL/min, or ibandronate if CrCl <30 mL/min • Annual Vitamin D level (25-hyroxyvitamin D) ng/mL (Zoledronic Acid, Ibandronate)			
Maintenance labs required: • Serum Creatinine (q3 months for ibandronate, q6 months for denosumab and romosozumab-aqqg, q12 months for zoledronic acid) • Serum Calcium every 6 months for denosumab, ibandronate OR romosozumab-aqqg; every 12 months for zoledronic acid • Vitamin D level (25-dyroxyvitamin D) every 12 months (Zoledronic Acid, Ibandronate)			
☑ IV Access: Access and/or maintain IV site in accordance with MHS IV Therapy P&P: Peripheral IV Device Site Selection, Insertion, Maintenance, and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Changes and Removal.			
Treatment Regimen: □ Denosumab (Prolia) 60 mg SQ every 6 months x 1 year □ Zoledronic Acid (Reclast) 5 mg IV infusion over at least 15 minutes x 1 dose • Recommended to have patient hold furosemide or torsemide morning of infusion □ Ibandronate (Boniva) 3 mg IV push over 30 seconds every 3 months x 1 year • Recommended to have patient hold furosemide or torsemide morning of dose □ Romosozumab-aqqg (Evenity) 210 mg (administered as 2 injections) SQ every month x 12 doses • Contraindicated in patient with history of stroke or myocardial infarction within the preceding year			
✓ Vital Signs: Check vital signs prior to dose. Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)			
If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.): • Consult MultiCare hypersensitivity guideline for treatment management • Notify provider of reaction, assessment and need for futher orders			
Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.			
Was consent obtained: ☐ Yes ☐ No (if yes, please send DOCUMENTATION of consent with order)			
Provider Signature	Print Name	 Date	Time
Another brand of drug, identical in form and content, may be dispensed unless checked Orders expire in 12 months**			

Name:

MRN #:

CSN #:

Age / Sex and Gender:

Pre-printed Order **OSTEOPOROSIS TREATMENTS**

MultiCare 🕰

