| ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN | | | | | |
|--|--|---|--|---|---|
| Allergies/Reactions: | | Fax all infusions to: 833 Please mark the approp infusion center: | -380-8800 | Allenmore Infusion Center Auburn Infusion Center Gig Harbor Infusion Services Puyallup Infusion Center | DHEC Infusion Center North Spokane Infusion Center North Star Lodge Infusion Center |
| ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated. | | | | | |
| Darbepoetin and Epoetin or biosimilar | | | | | |
| Patient Name: Requested Date of Service:// | | | | | |
| Date of Birth: / Patient Phone Number: () 🗅 May leave message | | | | | |
| ICD -10 Code: | | | | | |
| Diagnosis: | Anemia of CKD | | D63.1 | | |
| | CKD Stage III CKD Stage IV CKD Stage V, NOT on dialys Other | | □ N18.3 □ N18.4 □ N18.5 □ | | |
| Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs and provide review of inclusion and exclusion criteria per Erythropoiesis-Stimulating agents for CKD Guideline **If required documentation not received with order, scheduling of treatment will be delayed until complete information is available** | | | | | |
| Patient weight = lb/kg (required) | | | | | |
| Exclusion Criteria: Uncontrolled hypertension, active bleeding, erythropoietin-stimulating agent (ESA) resistance due to antibody development, or allergy to darbepoetin alfa. *Refer to pharmacy ESA guideline for further details. Pharmacy Consult for ESA Guideline Initiation dose = 0.45 mcg/kg SUBQ every 4 weeks (dose will be rounded to nearest syringe size per ESA guideline) Initiation dose for Epoetin alfa-epbx = 10,000 units SUBQ every 2 weeks Maintenance dose and frequency adjusted per pharmacy guideline Patient's BP will be monitored with each injection, if BP>180/100, if BP>180/100 hold ESA and contact provider for further instructions. Baseline Labs Required per protocol: Anemia labs (Ferritin, transferrin saturation (TSAT), Vitamin B-12, Folate, WNL as described in pharmacy protocol Serum Creatinine Hgb < 10.0 g/dl for initiation and drawn within 7 days of ESA dose Maintenance Labs: Hgb roir to dose (within 7 days of ESA dose) Iron labs (Ferritin, TSAT) every 3 months - pharmacist to order appropriate replacement per guideline Vitamin B-12 and Folate every 6 months - notify provider if not WNL V Access for patients needing iron replacement: Access and/or maintain IV site in accordance with MHS IV Therapy P&P Peripheral IV Device Site Selection, Insertion, Maintenance and Discontinuation; and Maintenance of Central Venous Catheters-Flushing, Dressing Change and Removal If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.) Consult MultiCare Hypersensitivity guideline for treatment/management Notify provider of reaction, assessment and need for further orders | | | | | |
| Provider Signature Print Name Date Time | | | | | |
| | ⁻ drug, identical in form and content, | | ss checked [| | |
| Patient Identification - Always Attach Patient Label Pre-printed Order | | | | | |
| Name: | | | DARBEPOETIN AND EPOTEIN OR BIOSIMILAR | | |
| MRN #: | | | | | |
| CSN #: | | | Mul | tiCare 🔁 | |
| Age / Sex and Gender: | | | | | |