ALL ORDERS MUST BE SIGNED, DATED AND TIMED BY PHYSICIAN			
Allergies/Reactions:	Fax all infusions to: 833-380-8800 Please mark the appropriate infusion center:	☐ Allenmore Infusion Center☐ Auburn Infusion Center☐ Gig Harbor Infusion Services☐ Puyallup Infusion Center☐	☐ DHEC Infusion Center☐ North Spokane Infusion Center☐ North Star Lodge Infusion Center☐
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), physicians are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.			
Ustekinumab (Stelara)			
Patient Name:	•	,	/ /
Patient Name: Requested Date of Service:/			
Diagnosis: ☐ Plaque psoriasis ☐ Psoriasis arthritis ☐ Crohn's disease ☐ Ulcerative colitis Required: H&P with documentation to support		code and supporting labs	
Required: H&P with documentation to support above diagnosis including ICD-10 code and supporting labs **If required documentation not received with order, scheduling of treatment will be delayed until complete information is available**			
☑ Reason patient not able to self-administer medication:			
Baseline labs required: • Latent TB testing Date://_ • HBV screening Date://_ • HCV screening Date://_ • HIV screening Date://_ • CBC, CMP	Results: Results:		
Maintenance labs required: • Annual Latent TB testing • CBC, CMP every 6 months			
Contact provider if patient develops serious infection; close monitoring by provider is recommended. Follow-up appointment with referring provider scheduled for (date):			
Patient weight = lb/kg (required)			
Treatment Regimen for Ustekinumaab (Stelara) dose: Plaque psoriasis: □ 45 mg SubQ (Wt <100 kg) □ 90 mg SubQ (Wt ≥100 kg) Psoriasis arthritis: □ 45 mg SubQ □ 90 mg SubQ (for pts w/ co-existent moderate to severe plaque psoriasis) Administer both at week 0 and 4 weeks later, followed by administration every 12 weeks			
Crohn's/Ulcerative Colitis: ☐ 260 mg IV over 1 hr x 1 dose (=55 kg) ☐ 390 mg IV over 1 hr x 1 dose (55-85 kg) ☐ 520 mg IV over 1 hr x 1 dose (>85 kg) Followed by 90 mg SUBQ 8 weeks after initial IV dose then every 8 weeks			
☑ Vital Signs: Check vital signs prior to and at completion of dose. Contact provider if systolic BP >180; diastolic BP >100; systolic BP <90; HR >110; temp >38C (100.4F)			
 If hypersensitivity develops (fever, chills, hypotension, rigors, itching, rash, etc.): Consult MultiCare hypersensitivity guideline for treatment management Notify provider of reaction, assessment and need for futher orders 			
Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.			
Was consent obtained: ☐ Yes ☐ No (if yes, please send DOCUMENTATION of consent with order)			
Provider Signature	Print Name	Date	Time
Another brand of drug, identical in form and content, may be dispensed unless checked Orders expire in 12 months**			
Patient Identification - Always Attach Patient Label Pre-printed Order			
Name:		EKINUMAB (St	elara)
AADNI #	031		ciai aj

Name:

MRN #:

CSN #:

Age / Sex and Gender:

MultiCare 🕰

