MHS Home/Alternate	Site Infusion Services	
Benralizumab (Fasenra) Injection Order Set		
ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER		
Patient Name:	DOB: /	
Patient Phone Number: ()	Requested Date of Service:/	
Patient Allergies:		
Diagnosis: ICD – 10 Code:		
Baseline Labs (Required): Absolute eosinophilic count > 0.015 K/uL in prior 6 weeks OR absolute eosinophilic count > 0.03 K/uL in prior 12 months. Date:/ Result:		
Lab Orders:		
Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. ** Emergency phone number for provider (required) **		
Provider Order for Benralizumab (Fasenra) Injection		
	se with check boxes), providers are responsible for indicating a ne order. Orders left unchecked will not be initiated.	
Biologic Injection: Benralizumab (Fasenra) Dosing: 30 mg SUBQ every 4 weeks for initial 3 doses followed by 40 mg SUBQ every 8 weeks 30 mg SUBQ every 8 weeks Skilled Nurse Interventions: Admit (first visit) patient to services for home infusion therapy of Benralizumab (Fasenra). Complete Skilled Nurse visit with each injection for ongoing home infusion therapy of Benralizumab (Fasenra). Obtain vital signs (TPR & B/P) at baseline and 30 minutes post-injection. Obtain patient weight at each visit. Draw labs as ordered. Inject Benralizumab [Fasenra] as prescribed. If vital signs stable 30 minutes after injection, Skilled Nurse will complete visit. If no injection-related events with previous 3 doses, may waive post-injection monitoring period and discharge pt home after completion.		
FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY		
Patient Identification: Name: MRN: DOB:	Pre-printed order – Page 1 of 2 Benralizumab (Fasenra) Order Set MultiCare	

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.): The Skilled Nurse will: Administer emergency medications as prescribed (below).

- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- Diphenhydramine 50 mg/mL solution: Inject 0.5 mL (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available. Max dose = 50 mg.
- Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2mL solution: Inject 2 mL (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.

101F/38.3C) and/or mild-to-moderate hypersensitivity red	vlaxis AND call 911 and provider. 50 mg) by mouth as needed for fever (PRN for temperature >
	a to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use
Code Status: Please note, patients will be considered FULL Code advance directive or living will, please include a copy with the o	
Was consent obtained: \square Yes \square No (if yes, please send DOCU	MENTATION of consent with order)
I certify that this patient is safe and appropriate to receive therap	y from Home Infusion Services.
Provider Signature: Printed	d Name: Date:
NPI: Orders expire in 12 months	s unless otherwise specified:
Provider/Clinic Information:	Return completed orders to: MultiCare Home/Alternate Site Infusion Services
Address:	253-459-6650 (phone) / 253-864-2785 (fax)
Phone #: Fax#:	
FOR MHS HOME/ALTERNATE SITE	INFUSION SERVICES USE ONLY
Patient Identification: Name:	Pre-printed order – Page 2 of 2 Benralizumab (Fasenra) Order Set
MRN: DOB:	MultiCare Revised 10/24