

MHS Home/Alternate Site Infusion Services Certolizumab Pegol (Cimzia) Injection Order Set

ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER

Patient Name: _____ **DOB:** ____/____/____ **Weight:** _____ lb/kg

Patient Phone Number: (____) ____-____ **Requested Date of Service:** ____/____/____

Patient Allergies: _____

Diagnosis: _____ **ICD – 10 Code:** _____

- Z45.2: Encounter for adjustment and management of vascular access device
- Z95.828: Presence of other vascular implants and grafts

Baseline Labs required:

- | | | |
|---------------------|----------------------|---------------|
| • CBC/CMP | Date: ____/____/____ | Result: _____ |
| • Latent TB testing | Date: ____/____/____ | Result: _____ |
| • HBV screening | Date: ____/____/____ | Result: _____ |
| • HCV screening | Date: ____/____/____ | Result: _____ |
| • HIV screening | Date: ____/____/____ | Result: _____ |

Reason patient is not able to self-administer medication: _____

Maintenance Labs required:

- CBC every 6 months
- Latent TB testing every 12 months

Lab Orders: _____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **** Emergency phone number for provider _____ (required) ****

Provider Order for Certolizumab Pegol (Cimzia) Injection

| | |
|-------------------------|--|
| ORDERS WITH CHECK BOXES | When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated. |
|-------------------------|--|

Biologic Injection: Certolizumab (Cimzia) Dosing per indication:

Ankylosing Spondylitis; Non-Radiographic Axial Spondyloarthritis; Psoriatic Arthritis, active; Rheumatoid Arthritis, active:

- Initial: 400 mg SUBQ at weeks 0, 2, and 4
 Maintenance:
 200 mg SUBQ every 2 weeks OR 400 mg SUBQ every 4 weeks

Crohn's disease, active:

- Initial: 400 mg SUBQ at weeks 0, 2, and 4
 Maintenance: 400 mg SUBQ every 4 weeks

Plaque Psoriasis:

- 400 mg SUBQ every 2 weeks OR
 For patients ≤ 90 kg 400 mg SUBQ at weeks 0, 2, and 4, then 200 mg SUBQ every 2 weeks

CONTINUED ON NEXT PAGE

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

| | |
|-----------------------|---|
| Name: MRN: DOB: | Pre-printed order – Page 1 of 2 Certolizumab Pegol (Cimzia) Order Set MultiCare |
|-----------------------|---|

Skilled Nurse Interventions:

- Admit (first visit) patient to services for home infusion therapy of Certolizumab Pegol (Cimzia).
- Complete Skilled Nurse visit with each injection for ongoing home infusion therapy of Certolizumab Pegol (Cimzia).
- Obtain vital signs (TPR & B/P) at baseline and post-injection.
- Obtain patient weight at each visit.
- Draw labs as ordered.
- Inject Certolizumab Pegol (Cimzia) as prescribed.
- If patient remains stable after the injection, Skilled Nurse will complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

- Administer emergency medications as prescribed (below).
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available. Max dose = 50 mg.
- **Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution:** Inject **2 mL** (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- **Acetaminophen (Tylenol) 325 mg tab:** Take **2 tablets** (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- **0.9% NaCl (NS) solution:** Infuse **500 mL** into the vein via gravity as needed at a rate needed to maintain IV access.
- **Oxygen:** Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.

Provider Signature: _____ Printed Name: _____ Date: _____

NPI: _____ Orders expire in 12 months unless otherwise specified: _____

Provider/Clinic Information:

Address: _____

Phone #: _____ Fax#: _____

Return completed orders to:

MultiCare Home/Alternate Site Infusion Services
253-459-6650 (phone) / 253-864-2785 (fax)

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name: _____
MRN: _____
DOB: _____

Pre-printed order – Page 2 of 2
Certolizumab Pegol (Cimzia) Order Set
MultiCare
Revised 10/24