MHS Home/Alternate Site Infusion Services		
Golimumab (Simponi Aria) Infusion Order Set ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER		
Patient Name:	DOB:/ Weight: lb/kg	
Patient Phone Number: ()	Requested Date of Service:/	
Patient Allergies:	requested bate of service.	
Diagnosis: ICD - 10 Code:		
 Z45.2: Encounter for adjustment and management of vascular access device Z95.828: Presence of other vascular implants and grafts 		
Baseline Labs Required: • Latent TB testing/Quantiferon: Date:/	Result: Result:	
Latent TB testing every 12 months CBC every 12 months		
Lab Orders:		
Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. ** Emergency phone number for provider (required) **		
Provider Order for Golimumab (Simponi Aria) Infusion		
·	, ,	
ORDERS WITH CHECK BOXES When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated. Biologic Infusion: Golimumab (Simponi Aria) administered in 100 mL NS infused over 30 minutes using 0.22 micron filter. Patient Weight:kg Initiation: 2 mg/kg = mg IV every 4 weeks for 2 doses, then every 8 weeks Maintenance: 2 mg/kg = mg IV every 8 weeks Additional Medications: O.9% NaCI (NS) PF flush in 10 mL syringe: Administer 5 – 30 mL IV as needed for establishing and maintaining IV access and for flushing IV line post infusion. Heparin: 100 units/mL for port OR 10 units/mL for other central line/midline, 3-5 mL PRN for de-accessing line. Cathflo (alteplase) 2 mg: Instill 2 mg (2 mL) into occluded catheter as needed for CVC/CVAD occlusion. Diphenhydramine 25 mg IV once, as needed for itching. Acetaminophen 325-650 mg tab/cap by mouth once, as needed for pain.		
FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY		
Patient Identification: Name: MRN: DOB:	Pre-printed order – Page 1 of 2 Golimumab (Simponi Aria) Order Set MultiCare	

Skilled Nurse Interventions:

- Admit (first visit) patient to services for home infusion therapy of Golimumab (Simponi Aria).
- Complete Skilled Nurse visit with each infusion for ongoing home infusion therapy of Golimumab (Simponi Aria).
- Obtain vital signs (TPR & B/P) at baseline and at completion of infusion.
- Obtain patient weight at each visit.
- Establish IV access and flush per policy to maintain patency.
- Draw labs as ordered.
- Withdraw ordered dose of Golimumab (Simponi Aria) (concentration prior to dilution is 12.5 mg/mL).
- Add reconstituted medication to prepared bag of Normal Saline 0.9% for a final volume of 100 mL (remove same volume of NS prior to adding medication).
- Infuse Golimumab (Simponi Aria) as prescribed.
- Once infusion is complete, flush IV line with Normal Saline 0.9% as prescribed.
- If vital signs are stable after infusion, Skilled Nurse will discontinue IV access and complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

DOB:

- STOP THE INFUSION.
- Administer emergency medications as prescribed (below).
- For mild infusion reaction, may restart the infusion at a slower rate (verified with pharmacist) if symptoms resolve with medication.
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access available. Max dose = 50 mg.
- Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution: Inject 2 mL (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- Acetaminophen (Tylenol) 325 mg tab: Take 2 tablets (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- 0.9% NaCl (NS) solution: Infuse 500 mL into the vein via gravity as needed at a rate needed to maintain IV access.
- Oxygen: Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders. Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)			
I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.			
Provider Signature: Printed	d Name:	_ Date:	
NPI: Orders expire in 12 months unless otherwise specified:			
Provider/Clinic Information:	Return completed orders to:		
Address:	MultiCare Home/Alternate Site Infusio 253-459-6650 (phone) / 253-864-278		
Phone #: Fax#:			
FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY			
Patient Identification: Name: MRN:	Pre-printed order – Page 2 of 2 Golimumab (Simponi Aria) Order Set MultiCare	Revised 10/24	