

MHS Home/Alternate Site Infusion Services Inclisiran (Leqvio) Injection Order Set

ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER

Patient Name:	DOB: ____/____/____	Weight: _____ lb/kg
---------------	---------------------	---------------------

Patient Phone Number: (____) ____-____	Requested Date of Service: ____/____/____
--	---

Patient Allergies: _____

Diagnosis: _____ ICD – 10 Code: _____

Baseline labs required:

Lipid profile (fasting or non-fasting) Date: ____/____/____ Result: _____

Maintenance labs required:

Lipid profile (fasting or non-fasting) 4-12 weeks after starting therapy
Lipid profile (fasting or non-fasting) every 3-12 months

Lab Orders: _____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **** Emergency phone number for provider _____ (required) ****

Provider Order for Inclisiran (Leqvio) Injection

ORDERS WITH CHECK BOXES	When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.
-------------------------	--

Medication:

- Initiation: Inclisiran (Leqvio) 284 mg SUBQ x 1. Repeat dose in 3 months (12 weeks), then continue every 6 months (24 weeks)
- Maintenance: Inclisiran (Leqvio) 284 mg SUBQ every 6 months (24 weeks)

Skilled Nurse Interventions:

- Admit (first visit) patient to services for home/alternate site injection therapy of Inclisiran (Leqvio).
- Complete Skilled Nurse visit with each injection for ongoing home/alternate site injection therapy of Inclisiran (Leqvio).
- Obtain vital signs (TPR & B/P) at baseline and at completion of injection.
- Obtain patient weight at each visit.
- Inject Inclisiran (Leqvio) as ordered.
- If patient is stable after injection and any observation period, Skilled Nurse will complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

- Administer emergency medications as prescribed (below).
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number for additional instructions.
- Notify MHS Home/Alternate Site Infusion Pharmacist AND Nurse Supervisor.

CONTINUED ON NEXT PAGE

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:
MRN:
DOB:

Pre-printed order – Page 1 of 2
Inclisiran (Leqvio)
MultiCare

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access available. Max dose = 50 mg.
- **Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution:** Inject **2 mL** (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) intramuscular every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- **Acetaminophen (Tylenol) 325 mg tab:** Take **2 tablets** (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- **0.9% NaCl (NS) solution:** Infuse **500 mL** into the vein via gravity as needed at a rate needed to maintain IV access.
- **Oxygen:** Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.

Provider Signature: _____ Printed Name: _____ Date: _____

NPI: _____ Orders expire in 12 months unless otherwise specified: _____

Provider/Clinic Information:

Address: _____

Phone #: _____ Fax#: _____

Return completed orders to:

MultiCare Home/Alternate Site Infusion Services
253-459-6650 (phone) / 253-864-2785 (fax)

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:

MRN:

DOB:

Pre-printed order – Page 2 of 2

Inclisiran (Leqvio)

MultiCare

Revised 12/24