

MHS Home/Alternate Site Infusion Services Ocrelizumab (Ocrevus) Infusion Order Set

ALL ORDERS MUST BE SIGNED AND DATED BY PRESCRIBING PROVIDER

Patient Name: _____	DOB: ____/____/____	Weight: _____ lb/kg
----------------------------	----------------------------	----------------------------

Patient Phone Number: (____) _____-_____	Requested Date of Service: ____/____/____
---	--

Patient Allergies: _____

Diagnosis: _____ **ICD – 10 Code:** _____

- Z45.2: Encounter for adjustment and management of vascular access device
- Z95.828: Presence of other vascular implants and grafts

Baseline Labs required:

- | | | |
|-----------------------|----------------------|---------------|
| • CBC and CMP | Date: ____/____/____ | Result: _____ |
| • HBV screening | Date: ____/____/____ | Result: _____ |
| • Latent TB screening | Date: ____/____/____ | Result: _____ |

Maintenance Labs required:

CBC and CMP prior to each infusion

Lab Orders: _____

Additional Requirements: In addition to order, please include H&P with documentation to support diagnosis. Must include ICD-10 Code(s) and supporting labs. **** Emergency phone number for provider _____ (required) ****

Provider Order for Ocrelizumab (Ocrevus) Infusion

ORDERS WITH CHECK BOXES	When an order is optional (those with check boxes), providers are responsible for indicating a check mark in the box next to the order. Orders left unchecked will not be initiated.
-------------------------	--

Pre-Medication(s):

- Methylprednisolone: 125 mg IV once, 30-60 minutes prior to infusion
- Acetaminophen: 650 mg tab/cap by mouth once, 30-60 minutes prior to infusion (patient may provide own supply)
- Diphenhydramine: 25 mg IV once, 30-60 minutes prior to infusion

Biologic Infusion:

Ocrelizumab (Ocrevus) added to 250 mL (300 mg) or 500 mL (600 mg) NS

- Initiation: 300 mg Day 1 and 15, then 600 mg every 6 months
- Maintenance: 600 mg every 6 months

Infuse _____mg via infusion pump using rate titration with 0.22 micron in-line filter

300 mg: begin at 30 mL/hr; increase by 30 mL/hr every 30 minutes to a max rate of 180 mL/hr (2.5 hrs or longer)

600 mg: begin at 40 mL/hr; increase by 40 mL/hr every 30 minutes to a max rate of 200 mL/hr (3.5 hrs or longer)

If no prior serious infusion reaction with any dose, may initiate rapid infusion starting with the 2nd maintenance dose:

100 mL/hr for 15 min, then 200 mL/hr for 15 min, then 250 mL/hr for 30 min, then 300 mL/hr for remaining infusion (2 hrs or longer). Pharmacist will adjust infusion rate based on patient response.

Additional Medications:

- 0.9% NaCl (NS) PF flush in 10 mL syringe: Administer 5 – 30 mL IV as needed for establishing and maintaining IV access and for flushing IV line post infusion.
- Heparin: 100 units/mL for port OR 10 units/mL for other central line/midline, 3-5 mL PRN for de-accessing line.
- Cathflo (alteplase) 2 mg: Instill 2 mg (2 mL) into occluded catheter as needed for CVC/CVAD occlusion.
- Diphenhydramine 25 mg IV once, as needed for itching.
- Acetaminophen 325-650 mg tab/cap by mouth once, as needed for pain.

CONTINUED ON NEXT PAGE

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:
MRN:
DOB:

Pre-printed order – Page 1 of 2
Ocrelizumab (Ocrevus) Order Set
MultiCare

Skilled Nurse Interventions:

- Admit (first visit) patient to services for home infusion therapy of Ocrelizumab (Ocrevus).
- Complete Skilled Nurse visit with each infusion for ongoing home infusion therapy of Ocrelizumab (Ocrevus).
- Obtain vital signs (TPR & B/P) at baseline, with each infusion rate titration, completion of infusion, and 60 minutes post-infusion.
- Obtain patient weight at each visit.
- Establish IV access and flush per policy to maintain patency.
- Draw labs as ordered.
- Administer pre-medications as prescribed.
- Add Ocrelizumab (Ocrevus) to 250 mL (300 mg) or 500 mL (600 mg) of sodium chloride 0.9% (NS).
- Infuse Ocrelizumab (Ocrevus) as prescribed.
- Once infusion complete, flush IV line with sodium chloride 0.9% (NS) as prescribed.
- If vital signs are stable 60 minutes after infusion, Skilled Nurse will discontinue IV access and complete visit.

If Hypersensitivity Reaction Develops (fever, chills, hypotension, rigors, itching, rash, etc.):

The Skilled Nurse will:

- STOP THE INFUSION.
- Administer emergency medications as prescribed (below).
- For mild infusion reaction, may restart the infusion at a slower rate (verified with pharmacist) if symptoms resolve with medication.
- Contact Emergency Medical Services (EMS/911) if indicated.
- Increase vital sign monitoring to every 5 minutes.
- Contact provider via emergency phone number _____ (required) for additional instructions.
- Notify MHS Home/Alternate Infusion Site Pharmacist AND Nurse Supervisor

Emergency Medications: To be administered by skilled nurse as needed for hypersensitivity reactions.

- **Diphenhydramine 50 mg/mL solution:** Inject **0.5 mL** (25 mg) into the vein via slow IV push over at least 2 minutes as needed for rash, itching, hives, flushing of skin, and/or mild-to-moderate hypersensitivity reaction. May give IM if no IV access is available. Max dose = 50 mg.
- **Methylprednisolone sodium succinate (SOLU-MEDROL) 125 mg/2 mL solution:** Inject **2 mL** (125 mg) into the vein as a slow IV push over at least 3 minutes once, as needed for hives, flushing of the skin, shortness of breath, chest tightness, tachycardia (HR>140 bpm), and/or moderate-to-severe hypersensitivity reaction. May give IM if no IV access is available.
- **EPINEPHrine (Adrenalin) 1 mg/mL solution:** Inject **0.3 mL** (0.3 mg) every 5-15 minutes as needed (maximum 3 doses) for severe allergic reaction or anaphylaxis AND call 911 and provider.
- **Acetaminophen (Tylenol) 325 mg tab:** Take **2 tablets** (650 mg) by mouth as needed for fever (PRN for temperature > 101F/38.3C) and/or mild-to-moderate hypersensitivity reaction.
- **0.9% NaCl (NS) solution:** Infuse **500 mL** into the vein via gravity as needed at a rate needed to maintain IV access.
- **Oxygen:** Initiate oxygen by RN PRN chest pain or dyspnea to keep SpO2>90%. For 1-6 LPM, use NC. For 6-10 LPM, use simple mask.

Code Status: Please note, patients will be considered FULL Code unless marked otherwise. If the patient has a POLST, advance directive or living will, please include a copy with the orders.

Was consent obtained: Yes No (if yes, please send DOCUMENTATION of consent with order)

I certify that this patient is safe and appropriate to receive therapy from Home Infusion Services.

Provider Signature: _____ **Printed Name:** _____ **Date:** _____

NPI: _____ Orders expire in 12 months unless otherwise specified: _____

Provider/Clinic Information:

Address: _____

Phone #: _____ Fax#: _____

Return completed orders to:

MultiCare Home/Alternate Site Infusion Services
253-459-6650 (phone) / 253-864-2785 (fax)

FOR MHS HOME/ALTERNATE SITE INFUSION SERVICES USE ONLY

Patient Identification:

Name:
MRN:
DOB:

Pre-printed order – Page 2 of 2
Ocrelizumab (Ocrevus) Order Set
MultiCare

Revised 10/24